

1.50" Seal Area | 12.00" | 1.50" Seal Area | 12.00" Seal Area

Tack Seal 1.5" From Top Of Bag (+/- 1/8")

Tack Seal .25" (+/- 1/8")

3" Customer Seal Area

4" (+/- 1/8") Micro Vent Holes

MAXLAWN CRABGRASS PREVENTER
with LAWN FOOD 22-0-4
MAXIMIZE YOUR LAWN

PREVENTS CRABGRASS AND FEEDS IN ONE APPLICATION

SEASON-LONG CRABGRASS CONTROL

LONG-LASTING FEEDING

STIMULATES DEEP GREEN COLOR AND GROWTH

1ST APPLICATION EARLY SPRING

COVERS 5,000 Sq. Ft. NET WEIGHT 16 LBS. (7.26 KG)

ACTIVE INGREDIENT: dithiopyr, 3-(5-pyridinylmethyl)carbothioic acid, 2-(difluoromethyl)-4-(2-methylpropyl)-5-(trifluoromethyl)-3,5-dimethyl ester, 0.122%
OTHER INGREDIENTS: 4-(2-methylpropyl)-5-(trifluoromethyl)-3,5-dimethyl ester, 99.878%
TOTAL: 100.000%
EPA Reg. No. 57131-15 EPA Est. No. 57131-W-1
This product contains 0.02 pounds of the active ingredient dithiopyr per 16 pound bag.
KEEP OUT OF REACH OF CHILDREN CAUTION
See back of bag for Precautionary Statements and Directions for Use.

GUARANTEED ANALYSIS
TOTAL NITROGEN (N).....22.0%
22.0% Urea Nitrogen*
SOLUBLE POTASH (K₂O).....4.0%
IRON (Fe).....1.0%
Plant Foods Derived From: Urea, Poly Coated Sulfur Coated Urea and Murate of Potash, Iron derived from Iron Oxide Sulfate.
*4.4% slowly available urea nitrogen from poly coated sulfur coated urea

100% SATISFACTION GUARANTEE ON YOUR MONEY BACK

PRECAUTIONARY STATEMENTS
Hazard to Humans and Domestic Animals
CAUTION: Causes moderate eye irritation. Avoid contact with eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

IF IN EYES Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

Net Use Number: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-424-9300 day or night, for emergency treatment information.

ENVIRONMENTAL HAZARDS
This product is toxic to fish and highly toxic to other aquatic organisms including oysters and shrimp. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn will help to prevent runoff to water bodies or drainage systems.

STORAGE AND DISPOSAL
Do not contaminate water, foodstuffs, feed or seed by storage or disposal. Pesticide Storage: Store this product only in its original container in a cool, dry area inaccessible to children and pets. For bags offer partial use, reseal opened bag by folding top down and securing. Pesticide Disposal: Wastes resulting from the use of this product that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticide disposal or in accordance with applicable federal, state or local procedures. **Container Handling:** Non-refillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling, if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

MAXLAWN Maximize Your Lawn Crabgrass Preventer with Lawn Food 22-0-4 provides a full feeding of fertilizer, pre-emergence control of crabgrass, and will control or suppress broadleaf weeds and grasses when applied prior to germination. In addition, this product will control crabgrass that has germinated within the past 4 weeks when crabgrass seedlings first become visible in the established lawn. Post-emergence control is limited to crabgrass. This product provides up to 4 months of crabgrass control when applied according to label directions.

WHERE TO APPLY
MAXLAWN Maximize Your Lawn Crabgrass Preventer with Lawn Food 22-0-4 should be applied only to lawns that are well established and to grass species listed on this label. The following grass species are tolerant to this product when applied as directed:

| Cool-Season Grasses | Warm-Season Grasses |
|--|---|
| Bermuda, creeping Bermuda, Kentucky Fescue, fine | Fescue, tall perennial Bahia Bermudagrass Buffalograss Cynodon Centipedegrass St. Augustinegrass Zoysia |

Note: For newly established lawns, initial application of this product should be made only after turfgrasses have developed a vigorous root system and received at least two mowings following seeding, sodding, or sprigging. Exception: In newly established lawns of Kentucky bluegrass, perennial ryegrass and/or tall fescue, this product may be applied two weeks after the first sign of germination.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

WHEN TO APPLY
Apply this product within a few days after mowing and delay mowing again for a few days after application. Do not apply this product in moderate or high wind conditions to prevent drift and uneven distribution of fertilizer. Avoid applying when temperatures are above 90°F.

Spring Application
Control of Crabgrass: This product provides pre-emergence control of crabgrass when applied prior to germination. In addition, this product provides post-emergence control of seedling crabgrass after germination and emergence. Post-emergence control is limited to crabgrass only and is effective up to 4 weeks after germination of crabgrass. This generally corresponds to the time when crabgrass seedlings first become visible in the established lawn. Properly timed applications will control existing crabgrass and prevent further emergence of crabgrass seedlings. Do not apply this product later than 4 weeks after crabgrass germination.

Control or Suppression of Other Grass and Broadleaf Weeds: For control or suppression of other grass and broadleaf weeds, this product must be applied as a pre-emergence treatment prior to their germination and emergence. Only crabgrass will be controlled with applications made up to 4 weeks after weed germination.

Fall Application
This product can also be applied from late summer through early fall (late August through November). When applied in the fall, a sequential or second application of this product must be made the following spring to provide season-long control of crabgrass. If this product is applied in the fall, do not reseed, overseed or sprig lawn within 12 weeks after application.

Weeds Controlled or Suppressed

| Grasses | Broadleaf Weeds |
|---|--|
| barley barnyardgrass crabgrass, annual brome crabgrass, large crabgrass, smooth crabgrass, Southern crabgrass cynodon dallisgrass goosegrass foxtail, green foxtail, yellow pigweed oats, wild ryegrass (annual & perennial) sarcobata stachys | bittercress † carpetweed chickweed † geranium, Carolina † henbit knockweed, prostrate lamb's ear † matricaria medic, black marestail oxalis, buttercup pineappleweed pigweed, redroot parsley plant † purslane, common rocket, London shepherdspurse † |

† Suppression only

Length of Control
MAXLAWN Maximize Your Lawn Crabgrass Preventer with Lawn Food 22-0-4 will provide up to 4 months of crabgrass control when applied according to label directions. The actual length of control will vary depending on use rate, weather conditions, condition of the lawn, intensity of weed pressure, and the location or "Region" within the United States where this product is being applied. Refer to the map below to determine the region of the U.S. that includes your location. Use the table below to determine the amount of this product to apply in your location based on the desired length of weed control or need for post-emergence crabgrass control.

Reseeding/Overseeding/Sprigging
Wait 10 weeks to reseed, overseed, or sprigging of areas treated with this product. When reseeding or overseeding, follow proper cultural practices for seedbed preparation, seeding, irrigation and fertilization.

Use "Regions" within the United States:

Use Rate:

| Region of U.S. | Amount to Apply (Pounds of product to cover per 1,000 square feet of lawn) | | |
|----------------|--|---------------------------|------------------------------|
| | 2-3 months | 3-4 months | Early Post-emergence Control |
| North | 2.4 lb./1,000 sq. ft. | 3.2 lb./1,000 sq. ft. | 4.0-9.2 lb./1,000 sq. ft. |
| South | 2.5 lb./1,000 sq. ft. | 4.0-4.7 lb./1,000 sq. ft. | 4.0-9.2 lb./1,000 sq. ft. |

Use the higher rate range in areas of high weed pressure or for increased length of control.

Maximum Application Rate

- Do not apply more than 9.2 lbs. of this product per 1,000 sq. ft. per application
- Do not apply more than 27.6 lbs. of this product per 1,000 sq. ft. per year
- In the state of New York, do not exceed 4.6 lbs. per 1,000 sq. ft. per year of this product

HOW TO APPLY
Apply evenly to the lawn using a drop or rotary-type spreader. Over-application may cause burning, under-application of this product will not achieve desired results. For best results, apply this product uniformly by walking or a steady speed carefully avoiding skipped areas. Calibrate the spreader according to the manufacturer's directions or follow Calibration Instructions below.

Watering or rainfall: Performance is improved if application is followed by 1/2 inch of rainfall or sprinkler irrigation. Erratic weed control may occur if not activated by rainfall or irrigation within 30 days of application.

SPREADING PATTERNS

BROADCAST SPREADER
First spread the borders of your lawn. Walk or an even speed making sure particles overlap previous pass (approximately within 5 feet of each other). Shut off spreader when stopped or when you reach a border strip.

| Brand/Model of Spreader | Calibration Settings at 3.2 lb./1,000 sq. ft. |
|---------------------------|---|
| AgriPro (B-30) | 12 |
| Chapin (B-30) | 1.5 |
| Chapin (B-15) | 4 |
| Earlway (E-30) | 12 |
| Earlway (E-30) (E-30) | 4.5 |
| Scotts (S-15) | 4.5 |
| Scotts (S-15) (S-15) | 4.5 |
| Scotts (S-15) Edge (E-15) | 2.5 |
| Scotts Easy Green | 6.5 |
| Scotts Speedy Green | 6.5 |

DROP SPREADER
First spread the borders of your lawn. Walk or an even speed making sure that spreader wheel overlaps previous wheel mark. Shut off spreader when stopped or when you reach a border strip.

| Brand/Model of Spreader | Calibration Settings at 3.2 lb./1,000 sq. ft. |
|-------------------------|---|
| AgriPro | 4 |
| Chapin | 4 |
| Earlway (E-30) (E-30) | 9.5 |
| LawnPro | 4 |
| Proline | 1.5 |
| Red Devil | 1.5 |
| Scotts (S-15) 1000 | 4 |
| Scotts (S-15) 2000 | 6 |
| Scotts (S-15) 3000 | 6 |
| Scotts (S-15) 4000 | 6.5 |

Calibration Instructions: Spreader Settings are recommendations. Variations in physical characteristics, walking speed, condition of spreader and application surface may require a different spreader setting. To provide proper distribution of product, while walking at a comfortable pace, check your spreader on 1,000 square feet of lawn and adjust accordingly to apply the 1,000 square foot rate.

Calculating Total Lawn Area
LENGTH X WIDTH = TOTAL SQ. FT. then subtract NON-LAWN AREAS (House, Garden, Driveway)

Warranty Disclaimer
Seller warrants that this product conforms to the chemical description of the label and is reasonably fit for the purpose stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. To the extent consistent with applicable law, SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

Terms and Conditions of Use
Read the entire Directions for Use, Conditions and Inherent Risks of Use before using this product. If terms are not acceptable, return the unopened product container at once. To the extent consistent with applicable law, by using this product, user or buyer accepts the following Conditions and Inherent Risks of Use.

CONDITIONS: The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product.

INHERENT RISKS OF USE: To the extent consistent with applicable law, the manufacturer, or the seller shall not be liable for consequential, special, or indirect damages resulting from the use, handling, application, storage, or disposal of this product or for damages in the nature of penalties, and the buyer and the user waive any right that they may have to such damages.

Barcode: 0 24378 97101 4
DPS # 57131-150-1516

DISTRIBUTED AND GUARANTEED BY EAU CLAIRE COOP OIL CO., PO. Box 837, Eau Claire, WI 54702

21.00" OD
20.50" ID

| COLORS | LPI | ANGLE | INK % |
|------------------|-------|-------|--------|
| LineBlack | Line | 0 | 4.36% |
| Black | 150 S | 38 | 12.24% |
| Cyan | 150 S | 8 | 6.58% |
| Magenta | 150 S | 68 | 5.42% |
| Yellow | 150 S | 83 | 6.42% |
| PANTONE Yellow C | Line | 0 | 23.37% |
| PANTONE 356 C | 65 C | 8 | 1.26% |

| JOB INFO | |
|-----------------|--|
| Customer: | EAU CLAIRE COOPERATIVE |
| Item Code #: | |
| Customer Job #: | |
| Dimensions: | 27.750" x 21.000" |
| Cylinder: | Z1* |
| Printer PDR: | |
| Printer Job#: | EAU010496 |
| Plate Material: | 067 DPR HDFTD |
| Orientation: | Reverse |
| Substrate: | Prints on Clear Poly-Lam to white poly |
| Artist: | cpooch@sgserver02 |
| Date: | 10/22/2019 |
| CSR: | LEAH TABORA |

| PROOF DETAILS | |
|---------------------------------|--|
| RO (09/21/2017) FIRST PROOF | |
| RO (09/21/2017) Edits to | |
| Spread Settings text. | |
| R1 -- 10.14.2019: Edits to text | |
| with supplied Illustrator file. | |
| R2 -- 10.17.2019: Edits to UPC | |

| CONTENT PROOF APPROVAL | |
|--|---|
| Description: | Max Lawn Crabgrass Preventer with Lawn Food 1st Application 22-0-4 16LB EAU010496 |
| Comments: | |
| CYBER GRAPHICS will not accept responsibility for any discrepancies in copy, legal requirements or design interpretation after final approval of this proof, unless we have not executed your requested changes. | |
| Client Approval | Printer Approval |
| [] NOT Approved for CONTENT | [] NOT Approved for CONTENT |
| [] Approved as is for CONTENT | [] Approved as is for CONTENT |
| Signature: _____ Date: _____ | Signature: _____ Date: _____ |

CYBERGRAPHICS
Perfecting packaging prepress. Protecting brands.

Design Code: EAU010496

NOVOLEX DIRECT PLASTICS

Artwork Approval

| APPROVED | APPROVED WITH |
|-------------------------------------|-------------------------------------|
| <input checked="" type="checkbox"/> | <input checked="" type="checkbox"/> |
| Copy & Content | |
| Package Size and Dimensions | |
| Inside Bag Dimensions | |
| Seal Area / No Print Area | |
| Number of Colours & PMS #'s | |
| Vent Holes and Positioning | |
| Eymark size and Positioning | |
| UPC Number | |

If colour square targets are shown on this proof they may appear on the finished product. They are required as part of our printing process however can be moved to a more appropriate location if requested.

Although every effort is made to produce the most accurate artwork possible, errors may occur. We do not assume any liabilities beyond the replacement of artwork.

PLEASE CHECK CAREFULLY!